

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12Q 1/04, G01N 33/02, 33/18</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/40980</b> <b>(43) International Publication Date:</b> 19 December 1996 (19.12.96)
<b>(21) International Application Number:</b> PCT/US96/08124 <b>(22) International Filing Date:</b> 31 May 1996 (31.05.96)  <b>(30) Priority Data:</b> 08/484,593      7 June 1995 (07.06.95)      US  <b>(71) Applicant:</b> IDEXX LABORATORIES, INC. [US/US]; One IDEXX Drive, Westbrook, ME 04092 (US).  <b>(72) Inventors:</b> TOWNSEND, David, E.; 29 Church Street, Scarborough, ME 04074 (US). CHEN, Chun-Ming; 17 Slocum Drive, Falmouth, ME 04105 (US).  <b>(74) Agents:</b> WARBURG, Richard, J. et al.; Lyon & Lyon, First Interstate World Center, Suite 4700, 633 West Fifth Street, Los Angeles, CA 90071-2066 (US).		<b>(81) Designated States:</b> AU, BR, CA, JP, MX, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> METHOD AND COMPOSITION FOR DETECTING BACTERIAL CONTAMINATION IN FOOD PRODUCTS		
<b>(57) Abstract</b> <p>This invention relates to a method for detecting the existence or measuring the concentration of total viable bacteria in a test sample from a food product. A medium is provided which contains three or more different enzyme substrates each having a nutrient moiety and a detectable moiety linked together. When a substrate is hydrolysed by a bacterial enzyme to create a separate detectable moiety, it causes or produces a detectable signal. These substrates produce detectable signals when any one of a phosphatase enzyme, a glycosidase enzyme or a peptidase enzyme is present in the medium.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

DESCRIPTIONMethod and Composition for Detecting  
Bacterial Contamination in Food ProductsField of the Invention

This invention relates to methods and compositions for detecting the existence or measuring the concentration of bacterial contamination in food products.

5 Background of the Invention

Ground beef and chicken are susceptible to rapid spoilage by psychotropic bacteria which thrive at refrigeration temperatures. As a result, these products have very short shelf-lives which are directly related to the  
10 initial concentration of contaminating bacteria.

Current methods for measuring the concentrations of bacterial contamination in ground beef and chicken include the standard plate count (Difco Laboratories) as well as the Petri Film system (3M) (see generally, Compendium of  
15 Methods for the Microbiological Examination of Foods, Third Edition, Edited by Carl Vanderzant and Don F. Splittstoesser, Compiled by the APHA Technical Committee on Microbiological Methods for Foods). These methods require around 48 hours of incubation in a 35°C incubator  
20 before the results can be read. Both methods utilize a solid nutrient base to support the growth of individual cells into bacterial colonies. Many food-borne bacteria are incapable of growing into colonies on these surfaces when incubated at 35°C; thus, the concentrations of total  
25 viable bacteria measured by the above methods may be underestimated.

In addition, the long incubation periods of these methods can cause these food products to remain in storage for several days until the concentrations of contaminating  
30 bacteria are known. If these tests could be completed in a shorter period of time it would allow companies to

release their products sooner so as to lower costs, increase sales, and provide better product to the consumer.

There have been attempts to measure the bacterial concentration in food by measuring specific metabolic by-products of individual microorganisms. These methods include: electrical impedance assays, ATP assays, antibody-based assays, and carbon-14 labelled substrate assays. Indicators of microbial growth have also been used to monitor the growth of target microbes which change color only after growth of the target microbe is detected. These indicators normally react chemically with a metabolic by-product produced by the target microbes resulting in a color change in the medium. Examples of chemicals which change color in the presence of pH changes associated with growth include phenol red, bromocresol blue, and neutral red. For example, Golber, U.S. Patent No. 3,206,317, uses phenol red, a chemical which changes color in the presence of acidic waste products produced by the target microbe. Berger et al., U.S. Patent No. 3,496,066, describes the use of compounds which bacteria convert to dyestuffs, e.g., tropinones and dioxanes, Bochner, U.S. Patent No. 4,129,483 describes using a non-biodegradable substance (tetrazolium) which is chemically reduced to produce a color change. In all of these examples, the indicator is a compound which does not serve as a source of a required nutrient.

Edberg (U.S. Patent No. 4,925,789), incorporated by reference herein, describes a selective growth medium for a microbe containing a nutrient indicator which can only be metabolized by a target microbe. When metabolized by a target microbe, the nutrient indicator releases a moiety which imparts a detectable change to the medium.

#### Summary of the Invention

The present invention relates to a bacterial growth medium and methods for detecting the existence or

measuring the concentration of bacteria in a test sample. The claimed medium and methods measure viable bacteria as a function of the activities of several classes of bacterial enzymes, including, but not limited to, phosphatases, glycosidases (such as glucosidases), and aminopeptidases. The presence of at least one of these groups of enzymes in any given bacterial species will be detected by the appearance of a detectable signal such as a fluorescent signal. Therefore, this invention is useful in detecting the existence or measuring the concentration of total viable bacteria or at least a multitude of viable bacteria in a test sample in a single assay. In specific examples, cocktails of enzyme substrates are made to measure the concentration of bacterial contamination in food products, such as ground beef and chicken.

Thus, in one aspect, the invention features a bacterial growth medium containing three or more different enzyme substrates each hydrolysed by a different bacterial enzyme to cause or produce a detectable signal.

In a preferred embodiment, the three or more different enzyme substrates each has both a nutrient moiety and a detectable moiety linked together by a covalent bond. Each of these enzyme substrates is hydrolysed by a different bacterial enzyme to produce a separate detectable moiety which causes or produces a detectable signal in the medium. In a further preferred embodiment, the detectable signals caused or produced are of identical type.

By "medium" is meant a solid, powder or liquid mixture which contains all or substantially all of the nutrients necessary to support bacterial growth. Amino acids, minerals, vitamins and other elements known to those skilled in the art to be necessary for bacterial growth are provided in the medium, including, but not limited to, those disclosed in U.S. application nos. 08/334,788 and 08/335,149, both filed on November 4, 1994, incorporated

by reference herein. In a preferred embodiment, the medium is liquid.

For example, the following components are provided in the medium in approximately the amounts indicated. Those  
5 in the art will understand that not every component is required. Components may also be substituted with other components of similar properties. The amounts of components may also be varied.

Amino acids may be provided from a variety of sources.  
10 These can be provided from natural sources (e.g., extracts of organisms), as mixtures, or in purified form. The natural mixtures may contain varying amounts of such amino acids and vitamins. Not all amino acids must be provided, and the relative amount of each can vary. For general  
15 guidance, specific amounts of such amino acids and vitamins are indicated below. These amounts are for guidance only and are not limiting in this invention. Those in the art will recognize that many different combinations of amino acids and vitamins can be used in  
20 the medium of this invention. The lists provided below exemplify just one such example. Normally, only amino acids which cannot be synthesized endogenously by the microorganisms to be detected must be provided. However, other amino acids may be provided without departing from  
25 the medium of the invention.

The medium preferably includes at least the following amino acids in approximately the following amounts (per liter of medium): Alanine (0.015 to 0.60 grams), Arginine (0.080 to 3.2 grams), Aspartic Acid (0.018 to 0.72 grams),  
30 Cystine (0.09 to 3.6 grams), Glutamic Acid (0.030 to 1.20 grams), Glycine (0.050 to 2.00 grams), Histidine (0.025 to 1.00 grams), Isoleucine (0.035 to 1.40 grams), Leucine (0.040 to 1.60 grams), Lysine (0.050 to 2.00 grams), Methionine (0.01 to 0.50 grams), Phenylalanine (0.01 to  
35 0.90 grams), Proline (0.02 to 2.80 grams), Serine (0.01 to 0.40 grams), Threonine (0.01 to 1.10 grams), Tryptophan

(0.002 to 0.26 grams), Tyrosine (0.01 to 1.20 grams), and Valine (0.02 to 1.10 grams).

Salts may be provided as a source of ions upon dissociation. Such salts may include (per liter of medium):  
5 potassium chloride (e.g., about 0.5 to 1.5 grams); copper sulfate (e.g., about 40 to 50  $\mu$ g); ammonium acetate or ammonium sulfate (e.g., about 4.0 to 6.0 grams); potassium iodide (e.g., about 50.0 to 150.0  $\mu$ g); ferric chloride (e.g., about 150.0 to 250.0  $\mu$ g); manganese sulfate (e.g.,  
10 about 300.0 to 500.0  $\mu$ g); sodium molybdate (e.g., about 150.0 to 250.0  $\mu$ g); zinc sulfate (e.g. about 300.0 to 500.0  $\mu$ g); and sodium chloride (e.g. about 0.05 to 0.15 g).

Other inorganic moieties may be included to aid microbial growth. These include the following (to the extent  
15 not already provided in the above sources of various chemical entities and described in amounts per liter): Phosphorus (about 0.5 mg), Potassium (about 0.4 mg), Sodium (about 30 to 60 mg), and trace amounts of Calcium,  
20 Magnesium, Aluminum, Barium, Chloride, Cobalt, Copper, Iron, Lead, Manganese, Sulfate, Sulfur, Tin and Zinc.

Vitamins required for growth and reproduction of the microorganism sought to be detected may also be provided. These can be provided in a pure form or as part of a more  
25 complex medium. Such vitamins may be present in approximately the following amounts (per liter of medium): Biotin (about 0.15 to 60  $\mu$ g), Pantothenic Acid (about 15.0 to 65.0  $\mu$ g), Pyridoxine (about 2.0 to 9.0  $\mu$ g), Riboflavin (about 10.0 to 50.0  $\mu$ g), Folic acid (about 5.00 to 50.00  
30  $\mu$ g), Thiamine (about 10.0 to 50.0  $\mu$ g), Vitamin B12 (about 0.20 to 0.50  $\mu$ g), and Niacin (about 15.0 to 55.0  $\mu$ g).

By "bacterial enzyme" is meant an enzyme whose enzymatic activity such as the ability to hydrolyse a substrate or a plurality of substrates is characteristic  
35 of a bacterium or a plurality of bacteria. In this invention, the enzymatic activities of a bacterial enzyme or bacterial enzymes are used to detect or measure the

concentration of bacteria in a test sample. The bacterial enzymes include all those known to one skilled in the art, including, but not limited to, those listed in Enzymes, 3rd edition, edited by Malcolm Dixon, Edwin C. Webb, C.J.R. Thorne, and K.F. Tipton, 1979, Academic Press, U.S.A. In a preferred embodiment, the bacterial enzyme is selected from the group consisting of alkaline phosphatase, acid phosphatase, esterase, lipase, N-acetyl- $\beta$ -D-galactosaminidase, N-acetyl- $\beta$ -D-glucosaminidase, Neuraminidase, L-arabinopyranosidase,  $\beta$ -D-fucosidase,  $\alpha$ -L-fucosidase,  $\beta$ -L-fucosidase,  $\alpha$ -D-galactosidase,  $\beta$ -D-galactosidase,  $\alpha$ -D-glucosidase,  $\beta$ -D-glucosidase,  $\beta$ -D-glucuronidase,  $\alpha$ -D-mannosidase, pyrophosphatase, sulfatase,  $\beta$ -D-xylosidase, peptidase (preferably an aminopeptidase, more preferably an (L or D amino acid) - aminopeptidase), trypsin, chymotrypsin, and phosphohydrolase.

By "substrate" is meant a molecule or substance on which a bacterial enzyme acts. The enzymatic reaction usually involves hydrolysing one or more covalent bonds, forming one or more covalent bonds, or both. A covalent bond in the substrate between the nutrient moiety and the detectable moiety is hydrolysed by a bacterial enzyme to produce a separate detectable moiety. The substrates include all those known to one skilled in the art, including, but not limited to, those in the product listing of AerChem, Inc. with detectable moieties attached thereto (see Table I).

By "nutrient moiety" is meant a molecule or substance which is a nutrient or metabolic source for a bacterium, including, but not limited to, vitamins, minerals (e.g., phosphorus in the form of phosphate), trace elements, amino acids (e.g., L-alanine), carbon (e.g., glucose), or nitrogen.

By "detectable signal" is meant a characteristic change in a medium or sample that is observable or measurable by physical, chemical, or biological means known to those skilled in the art. Such a detectable



signal may be a change in emission or absorbance of visible or invisible light or radio waves at a certain wavelength, electrical conductivity, hybridization, enzymatic reaction, emission of gas, or odor. A detectable signal may also be a change in physical state such as between solid, liquid and gas. In preferred embodiments, detectable signals include a change in color or fluorescent emission of the medium.

By "identical type of detectable signal" is meant that the separate detectable moieties hydrolysed from different enzyme substrates cause or produce detectable signals that are measurable by the same or substantially the same physical, chemical or biological parameter, including, but not limited to, color, fluorescent emission, odor, enzymatic reaction, hybridization, or electric conductivity (although the intensity or quantity of signals caused or produced by different separate detectable moieties may be different). For example, yellow colors of different intensity would be considered of the identical type. Color change and fluorescence would not be considered to be identical type of detectable signal.

By "detectable moiety" is meant a molecule or substance which can be covalently linked to a nutrient moiety or exists as a separate entity by itself. The detectable moiety does not cause or produce a detectable signal when it is covalently bonded to a nutrient moiety. However, when an enzyme from a bacterium hydrolyses the substrate, a detectable moiety is released and causes or produces a detectable signal. In preferred embodiments, the detectable moieties are chromogens which produce a color change observable in the visible wavelength range or fluoresces when properly excited by an external energy source. Examples of detectable moieties include, but are not limited to, orthonitrophenyl, phenolphthalein, and 4-methylumbelliferone moieties.

The invention also features a method of using the medium to detect the existence or measure the concen-

tration of bacterial contamination in a test sample. The medium is inoculated with the test sample and incubated under a condition suitable for bacterial growth for a certain time period (preferably no more than 24 hours, more preferably no more than 15 hrs, even more preferably no more than 10 hours). Then the detectable signal is measured as an indication of the concentration of bacteria in the test sample. Using this method, a detectable signal is produced when at least one of the three or more different bacterial enzymes is or are present in the bacteria which are incubating in the medium.

By "test sample" is meant a piece, fraction, aliquot, droplet, portion, fragment, volume, or tidbit taken from a food product such as ground beef or chicken, a human or animal test subject, a soil, water, air or other environmental source, or any other source whose bacterial concentration is to be measured. A test sample may be taken from a source using techniques known to one skilled in the art, including, but not limited to, those described or referred to in Compendium of Methods for the Microbiological Examination of Foods, Third Edition, Edited by Carl Vanderzant and Don F. Splittstoesser, Compiled by the APHA Technical Committee on Microbiological Methods for Foods, incorporated by reference herein.

By "bacteria" is meant one or more viable bacteria existing or co-existing collectively in a test sample. The term may refer to a single bacterium (e.g., *Aeromonas hydrophilia*, *Aeromonas caviae*, *Aeromonas sobria*, *Streptococcus uberis*, *Enterococcus faecium*, *Enterococcus faecalis*, *Bacillus sphaericus*, *Pseudomonas fluorescens*, *Pseudomonas putida*, *Serratia liquefaciens*, *Lactococcus lactis*, *Xanthomonas maltophilia*, *Staphylococcus simulans*, *Staphylococcus hominis*, *Streptococcus constellatus*, *Streptococcus anginosus*, *Escherichia coli*, *Staphylococcus aureus*, *Mycobacterium fortuitum*, and *Klebsiella pneumonia*), a genus of bacteria (e.g., streptococci, pseudo-

monas and enterococci), a number of related species of bacteria (e.g., coliforms), an even larger group of bacteria having a common characteristic (e.g., all gram-negative bacteria), a group of bacteria commonly found in a food product, an animal or human subject, or an environmental source, or a combination of two or more bacteria listed above. The bacteria include those described or referred to in Bergey's Manual of Systematic Bacteriology, 1989, Williams and Wilkins, U.S.A., incorporated by reference herein.

In preferred embodiments, one of the substrates is hydrolysed by the enzyme alkaline phosphatase; another substrate is hydrolysed by the enzyme glycosidase, including, but not limited to,  $\beta$ -D-glucosidase; and a third substrate is hydrolysed by a peptidase (preferably an aminopeptidase, more preferably an (L or D amino acid) - aminopeptidase), including, but not limited to, L-alanine aminopeptidase; the detectable moiety is a fluorescent moiety such that when the detectable moiety is hydrolysed from a substrate, it causes or produces a fluorescent signal; the medium contains at least the following three substrates: 4-methylumbelliferyl phosphate, 4-methylumbelliferyl- $\beta$ -D-glucoside and L-alanine-7-amido-4-methyl coumarin; and the medium is inoculated with a test sample from a food product, including, but not limited to, ground beef, chicken, milk, dairy products, and drinking water.

In another aspect, the invention features a bacterial growth medium containing two or more different enzyme substrates each hydrolysed by a different bacterial enzyme to cause or produce an identical type of detectable signal.

In a preferred embodiment, the two or more different substrates each has both a nutrient moiety and a detectable moiety linked together by a covalent bond. Each of these substrates is hydrolysed by a different bacterial

enzyme to produce a separate detectable moiety which causes or produces an identical type of detectable signal.

The invention also features a method of using the medium to detect the existence or measure the concentration of bacteria in a test sample. The medium is inoculated with the test sample and incubated, under a condition suitable for bacterial growth for a certain time period (preferably no more than 24 hours, more preferably no more than 15 hrs, even more preferably no more than 10 hours). Then the detectable signal is measured as an indication of the concentration of bacterial contamination in the test sample. Using this method, a detectable signal is produced when at least one of the two or more different bacterial enzymes is present in the incubation medium.

In preferred embodiments, the substrates are hydrolysed by an enzyme selected from the group consisting of alkaline phosphatase, glycosidase (which includes, but is not limited to,  $\beta$ -D-glucosidase), and peptidase (preferably an aminopeptidase, more preferably an (L or D amino acid) - aminopeptidase, including, but not limited to, L-alanine aminopeptidase); and the detectable moiety and the medium are analogous to those noted above.

In other embodiments, the invention uses the apparatus described by Naqui et al. in U.S. patent application 08/201,110, incorporated by reference herein, to quantify the concentration of bacterial contamination. An example of such an apparatus is sold by Idexx Laboratories Inc. under the name of Quanti Tray™. The quantifying step involves providing a test sample in a liquid form. The sample is placed or dispensed into the sample holding bag described by Naqui et al., and mixed with a medium to allow and promote growth of target bacteria within individual compartments. The mixture is incubated and the quantity and quality of the color or fluorescence change in each compartment is detected. The quantity and quality of positive compartment (i.e., a compartment having a detec-

table color or fluorescence change) is compared to a most probable number table which relates that value to the bacterial concentration of the test sample.

This invention has many advantages over the methods currently used to measure bacterial contamination. One advantage is its relatively short time to results. Certain psychotropic bacteria grow very slowly and can take from 48 to 72 hours before their colonies become large enough to count on an agar plate. However, count-  
10 able colonies need not be present for the results of Applicant's test to be read. The fluorescent color produced by these bacteria in the invention appears much faster than their corresponding colonies which results in a much shorter detection time. Applicant's test can  
15 reduce the incubation period to 24 hours or less.

Another advantage of the invention has over standard methods is the absence of interference by bacterial overgrowth. This is a particular problem when *Bacillus* species are present because they tend to grow over other  
20 bacterial colonies in such a way that the plate is unreadable. The *Bacillus* species are common in food, particularly those that have been heat treated, such as pasteurized milk. This problem is avoided in the invention because it does not depend on counting  
25 individual bacterial colonies.

This invention can be used in microbiology laboratories involved in end product testing and/or quality control of food products, the meat and poultry industries, the dairy industry, and the water industry. The invention  
30 may be used to measure the concentration of total viable bacteria in drinking water.

This invention also relates to a growth medium and methods for detecting or measuring the concentration of yeasts, fungi, or other eukaryotic microorganisms in a  
35 test sample using a formulated medium and steps like those described above.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments, and from the claims.

#### Description of the Preferred Embodiments

5 In the following description, reference will be made to various methodologies known to those of skill in the chemical, biological and microbiological arts. Publications and other materials setting forth such known methodologies to which reference is made are incorporated herein  
10 by reference in their entireties as though set forth in full. The compositions, methods, and products of this invention are applicable to biological and environmental specimens, and are useful in the chemical, biological and microbiological arts for the detection of bacterial conta-  
15 mination.

#### Detecting Bacteria by Measuring Bacterial Enzyme Activities

Bacteria derive their nutrients from an array of sources. The ability to metabolize certain sources may be  
20 characteristic of a particular bacterium or group of bacteria. Families, groups or species of bacteria may share enzyme specificity for certain nutrients which are lacking in other bacteria. By taking advantage of the metabolic characteristics of bacteria, it is possible to  
25 test for the presence of these enzyme systems, and thus, the bacteria which display these enzyme systems themselves. See Edberg, supra. Many enzymes have been identified which are specific to particular groups of bacteria and others likely will be identified in the  
30 future (see generally, Bergey's Manual of Systematic Bacteriology, 1989, Williams and Wilkins, U.S.A.).

For example, most gram negative bacteria, as a group, have L-alanine aminopeptidase enzyme activity. Substrates such as L-alanine- $\beta$ -orthonitrophenyl,  $\beta$ -naphthalamide- $\beta$ -L-  
35 alanine,  $\alpha$ -naphthol- $\beta$ -L-alanine, 4-methylumbelliferyl- $\beta$ -L-

alanine, and L-alanine-7-amido-4-methyl coumarin may be used in the medium to test for the presence of gram negative bacteria. The enzyme  $\beta$ -D-glucosidase is found in the *Enterococcus* group of bacteria. The enzyme may catalyze the hydrolysis of appropriate substrates containing chromogenic or fluorogenic moieties linked to a  $\beta$ -glucoside. This property may be used to indicate the presence or absence of enterococci in a sample. Substrates such as 4-methylumbelliferyl- $\beta$ -D-glucopyranoside may be used to indicate the presence of enterococci. *Staphylococcus aureus* is capable of hydrolysing orthonitrophenyl phosphate. Thus, if the growth medium contains this substrate as a source of phosphate, *Staphylococcus aureus* will grow and a color change will be produced by the release of the orthonitrophenyl moiety. *Mycobacterium fortuitum* requires  $\text{SO}_4$  as its source of sulfur, and this species can hydrolyse phenolphthalein-sulfate. Thus, in a selective medium whose only sulfur source is phenolphthalein-sulfate, this species will grow and produce a characteristic color change by release of the colored moiety. Furthermore, the enzyme  $\beta$ -D-glucuronidase is present in *E. coli*. Substrates such as orthonitrophenyl- $\beta$ -D-glucuronide,  $\beta$ -naphthalamide- $\beta$ -D-glucuronide,  $\alpha$ -naphthol- $\beta$ -D-glucuronide or methylumbelliferyl- $\beta$ -D-glucuronide may be used in a medium for the detection of *E. coli*.

#### Substrates and Detectable Moieties

Substrates including a chromogenic moiety have been demonstrated to display a characteristic color change in samples containing target bacteria having a bacterial enzyme capable of hydrolysing the substrates. For example, in the presence of  $\beta$ -D-glucuronidase, orthonitrophenyl- $\beta$ -D-glucuronide produces a color change to yellow, 4-methylumbelliferyl- $\beta$ -D-glucuronide produces fluorescence after excitation at 366 nm, and bromo-chloro-indole- $\beta$ -D-glucuronide produces a color change to blue when *E. coli* is present. In the presence of  $\beta$ -D-galactosidase, ortho-

nitrophenyl- $\beta$ -D-galactopyranoside produces a color change to yellow and 4-methylumbelliferyl- $\beta$ -D-galactopyranoside produces fluorescence after excitation at 366 nm when *E. coli* is present.

5 Two substrates producing different types of detectable signals have been used for detecting the presence of *E. coli* among total coliform bacteria. 4-methylumbelliferyl- $\beta$ -D-glucuronide may be used together with orthonitrophenyl- $\beta$ -D-galactopyranoside. If any *E. coli* is present, the  
10 sample solution both changes color to yellow and emits fluorescence after excitation at 366 nm.

Table I is a list of substrates from AerChem, Inc. that may be used to detect bacterial enzyme activities.

A detectable moiety may be attached to a nutrient  
15 moiety by methods known to those skilled in the art. The methods generally feature coupling or conjugating a nutrient moiety to a detectable moiety, such as a chromogenic moiety. Examples of such methods are described by Edberg in U.S. Patent Number 4,925,789, incorporated by  
20 reference herein.

The following non-limiting example features a liquid based bacterial growth medium used to quantify the total number of viable bacteria present in ground beef and chicken. This medium comprises 4-methylumbelliferyl  
25 phosphate (MUP), 4-methylumbelliferyl- $\beta$ -D-glucoside (MUD), and L-alanine-7-amido-4-methyl coumarin (ala-AMC). An example of the composition is described in Table II. The composition of defined media is described in Table III. MUP, MUD, ala-AMC, and potassium nitrate were purchased  
30 from Sigma. Bacto Proteose Peptone No. 3 was purchased from DIFCO.

The substrate 4-methylumbelliferyl- $\beta$ -D-glucoside is used to detect the presence of the enzyme  $\beta$ -D-glucosidase which is present in *Streptococci*, *Enterococci*, and other  
35 related bacteria commonly found in fresh meat.

The substrate L-alanine-7-amido-4-methylcoumarin is used to detect the presence of the enzyme L-alanine amino-



peptidase which is found in most *pseudomonas* species and other gram negative bacteria. Applicant discovered that this substrate is particularly sensitive to the presence of psychotropic bacteria which cause spoilage in meat.

5 Other substrates can be used in place of L-alanine-7-amido-4-methylcoumarin to detect other types of aminopeptidases in this group of bacteria without sacrificing sensitivity.

The substrate 4-methylumbelliferyl phosphate is used  
10 to detect the presence of phosphatases such as alkaline phosphatase and acid phosphatase which are found in most bacterial species. This enzyme substrate supports the detection of bacteria which lack or have diminished L-alanine aminopeptidase and  $\beta$ -D-glucosidase activities.

15 Because phosphatase,  $\beta$ -D-glucosidase, and L-alanine aminopeptidase are present in the vast majority of bacteria which contaminate ground beef and chicken, only one of these enzymes needs to be functional in the food-borne bacteria for viability to be detected. This test,  
20 therefore, has built-in redundant screens which support a highly accurate measure of total viable bacteria in ground beef and chicken.

The presence of bacteria is indicated by the appearance of a blue fluorescent color in the medium after it is  
25 exposed to an external ultra-violet lamp (366 nm wavelength). This test yields result after no more than 24 hours of incubation at 35°C.

The substrates MUP, MUD, or ala-AMC are hydrolysed by phosphatase,  $\beta$ -D-glucosidase, or L-alanine aminopeptidase  
30 to produce both nutrient and fluorescent moieties. The nutrient moieties (i.e., phosphate, glucose, and L-alanine) are consumed by the bacteria as a part of their normal metabolism. The fluorescent moieties (i.e., 4-methylumberiferone or 7-amino-4-methyl coumarin) produce  
35 fluorescent signals (maximum emission at 450 nm) which are used as indicators of bacterial viability.

The time required for the fluorescent color to appear is dependent upon the concentration of bacteria present in the reagent. Higher concentration of viable bacteria in the medium results in a proportional decrease in the time required for color development. Therefore, this test can be adapted to instrumentation because of the linear relationship between bacterial concentration and time to signal development, such as that described in Naqui et al., U.S. application no. 08/201,110, hereby incorporated by reference.

Naqui et al. describes an accurate method for quantifying the number of bacteria in a liquid sample. The invention employs a novel apparatus for holding a liquid sample. The apparatus features a bag which is designed for receiving a liquid sample and subsequently distributes the liquid sample into separate compartments within the bag so that different aliquots of one or more sizes may be tested. The invention described in that application further allows quantifying the microorganisms present in the sample by adding a medium to promote growth of microorganisms, heat sealing the bag of the invention for about five seconds at a temperature of about 250°F to 350°F, incubating the sample at an appropriate temperature for an appropriate length of time to allow growth of microorganisms, and recording and analyzing the results. The quantifying step involves detecting the quantity and quality of the color change in each compartment, and comparing that quantity and quality to a most probable number table which relates that value to the bacterial concentration of the test sample.

For example, each 10 ml Quanti Tray™ system contains 50 individual wells capable of holding 0.2 ml of medium. A 51st well is present which collects any "overflow" of medium not distributed into the first 50 wells. To begin the test the powder containing enzyme substrates is first dissolved in 10 ml of sterile water. Next, the reagent is inoculated with a predetermined volume of homogenized food

material. Finally, the reagent is sealed in a 10 ml Quanti Tray™ and placed in a 35°C incubator for 24 hours. The number of fluorescent wells present after incubation is compared against a most probable number (MPN) chart to  
5 determine the original concentration of bacteria present in the sample of food. Food containing higher than acceptable concentrations of contaminating bacteria can be retested to verify the results and/or disposed of to prevent distribution.

10 Because not all food is contaminated by the same bacteria found in ground beef and chicken, other enzyme targets may need to be selected to measure the total bacterial concentration of other types of food.

To design a medium for measuring the concentration of  
15 bacterial contamination in a test sample from another type of food or other sources prone to bacterial contamination, methods known to those skilled in the art (including, but not limited to, plating, nucleic acid hybridization study, microscopic observation, etc.) are used to identify bac-  
20 teria species existing in the sample. Once the bacteria species are identified, one skilled in the art would be able to identify an enzyme or a group of enzymes that are characteristic of the bacteria species, and substrates acted on by the enzymes. Substrates having a nutrient  
25 moiety and a detectable moiety linked together by a covalent bond that is hydrolysed by the enzymes are produced to be used in the medium.

All publications referenced are incorporated by reference herein, including the nucleic acid sequences and  
30 amino acid sequences listed in each publication. All the compounds disclosed and referred to in the publications mentioned above are incorporated by reference herein, including those compounds disclosed and referred to in articles cited by the publications mentioned above.

35 Other embodiments of this invention are disclosed in the following claims.

4-MU-SUBSTRATES (4-Methylumbelliferyl-Substrates)

- Bis(4-methylumbelliferyl)-phosphate  
 Bis(4-methylumbelliferyl)-phosphate Sodium salt  
 4-Methylumbelliferyl-acetate
- 5 4-Methylumbelliferyl-N-acetyl- $\beta$ -D-galactosaminide  
 4-Methylumbelliferyl-N-acetyl- $\beta$ -D-glucosaminide  
 2'-(4-Methylumbelliferyl)- $\alpha$ -D-N-acetyl-neuraminic acid Sodium salt  
 4-Methylumbelliferyl- $\alpha$ -L-arabinopyranoside  
 4-Methylumbelliferyl-butyrate
- 10 4-Methylumbelliferyl- $\beta$ -D-celloblopyranoside  
 4-Methylumbelliferyl- $\beta$ -D-celotriose  
 4-Methylumbelliferyl- $\beta$ -D-N,N'-diacetyl-chitobloside  
 4-Methylumbelliferyl-elaldate  
 4-Methylumbelliferyl- $\beta$ -D-fucoside
- 15 4-Methylumbelliferyl- $\alpha$ -L-fucoside  
 4-Methylumbelliferyl- $\beta$ -L-fucoside  
 4-Methylumbelliferyl- $\alpha$ -D-galactoside  
 4-Methylumbelliferyl- $\beta$ -D-galactoside  
 4-Methylumbelliferyl- $\beta$ -D-galactoside-6-phosphate Ammonium salt
- 20 4-Methylumbelliferyl- $\alpha$ -D-glucoside  
 4-Methylumbelliferyl- $\beta$ -D-glucoside  
 4-Methylumbelliferyl- $\beta$ -D-glucuronide  
 4-Methylumbelliferyl- $\alpha$ -guanidinobenzoate hydrochloride  
 4-Methylumbelliferyl-heptanoate
- 25 4-Methylumbelliferyl- $\alpha$ -L-iduronide  
 4-Methylumbelliferyl-laurate  
 4-Methylumbelliferyl-lignocerate  
 4-Methylumbelliferyl- $\alpha$ -D-mannoside  
 4-Methylumbelliferyl-nonaoate
- 30 4-Methylumbelliferyl-oleate  
 4-Methylumbelliferyl-palmitate  
 4-Methylumbelliferyl-phosphate (free acid)  
 4-Methylumbelliferyl-phosphate(di(2-amino-2-methyl-1,3-propanediol)salt  
 4-Methylumbelliferyl-phosphate Dicyclohexylammonium salt
- 35 4-Methylumbelliferyl-phosphate Disodium salt  
 4-Methylumbelliferyl-propionate  
 4-Methylumbelliferyl-pyrophosphate diester Disodium salt  
 4-Methylumbelliferyl-stearate  
 4-Methylumbelliferyl-sulfate Potassium salt
- 40 4-Methylumbelliferyl-6-sulfo-N-acetyl- $\beta$ -D-glucosaminide  
 4-Methylumbelliferyl- $\beta$ -D-N,N',N'-triacetylchitotriose  
 4-Methylumbelliferyl-4-trimethylammonium cinnamate chloride  
 4-Methylumbelliferyl- $\beta$ -D-xylose

AMC-SUBSTRATES (7-Amido-4-methylcoumarin-Substrates)

- N-a-Acetyl-lysine-7-amido-4-methylcoumarin acetate  
 N-Acetyl-L-phenylalanyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 L-Alanine-7-amido-4-methylcoumarin
- 5  $\beta$ -Alanine-7-amido-4-methylcoumarin TFA  
 D-Alanine-7-amido-4-methylcoumarin TFA  
 L-Alanine-4-amido-7-methylcoumarin TFA  
 L-Alanine-7-amido-4-methylcoumarin TFA  
 L-Alanine-7-amido-4-trifluoro-methylcoumarin TFA
- 10 L-Alanyl-L-alanyl-L-phenylalanine-7-amido-4-methylcoumarin  
 L-Alanyl-L-alanyl-L-phenylalanine-7-amido-4-methylcoumarin TFA  
 D-Alanyl-L-leucyl-L-lysine-7-amido-4-methylcoumarin  
 L-Alanyl-L-phenylalanyl-L-lysine-7-amido-4-methylcoumarin salt  
 L-Arginine-7-amido-4-methylcoumarin-hydrochloride
- 15 L-Arginyl-L-arginine-7-amido-4-methylcoumarin trihydrochloride  
 L-Asparagine-7-amido-4-methylcoumarine TFA  
 L-Aspartic acid- $\beta$ - (7-amido-4-methylcoumarin)  
 N-a-Benzoyl-DL-arginine-7-amido-4-methylcoumarin hydrochloride  
 N-a-Benzoyl-L-arginine-7-amido-4-methylcoumarin hydrochloride
- 20 N-Benzoyl-L-phenylalanyl-L-valyl-L-arginine-7-amido-4-methylcoumarin  
 hydrochloride  
 N-Benzoyl-L-valyl-glycyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 S-Benzyl-L-cysteine-7-amido-4-methylcoumarin  
 N-BOC-L-Phenylalanyl-L-seryl-L-arginine-7-amido-4-methylcoumarin acetate
- 25 N-BOC-L-Valyl-glycyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 N-BOC-L-Valyl-Ueucyl-L-lysine-7-amido-4-methylcoumarin Salt  
 N-a-CBZ-L-Arginine-7-amido-4-methylcoumarin hydrochloride  
 N-CBZ-Glycylglycyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 N-CBZ-Glycylglycyl-L-leucine-7-amido-4-methylcoumarin
- 30 N-CBZ-Glycyl-L-proline-7-amido-4-methylcoumarin  
 N-CBZ-Glycyl-L-prolyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 N- $\beta$ -CBX-L-Lysine-7-amido-4-methylcoumarin hydrochloride  
 N-CBZ-L-Phenylalanyl-L-arginine-7-amido-4-methylchloride hydrochloride  
 N-CBZ-L-Prolyl-L-arginine-7-amido-4-methylcoumarin hydrochloride
- 35 L-Citrulline-7-amido-4-methylcoumarin hydrobromide  
 L-Citrulline-7-amido-4-methylcoumarin TFA  
 D-Glutamic acid- $\gamma$ - (7-amido-4-methylcoumarin)  
 L-Glutamic acid- $\alpha$ - (7-amido-4-methylcoumarin)  
 L-Glutamine-7-amido-4-methylcoumarin hydrochloride
- 40 Glutaryl-glycyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 Glutaryl-glycylglycyl-L-phenylalanine-7-amido-4-methylcoumarin  
 Glutaryl-glycylglycyl-L-phenylalanine-7-amido-4-methylcoumarin  
 Glutaryl-L-phenylalanine-7-amido-4-methylcoumarin  
 Glycine-7-amido-4-methylcoumarin hydrobromide
- 45 Glycyl-L-alanine-7-amido-4-methylcoumarin hydrochloride  
 Glycyl-L-arginine-7-amido-4-methylcoumarin salt  
 Glycylglycine-7-amido-4-methylcoumarin hydrochloride

- Glycyl-L-phenylalanine-7-amido-4-methylcoumarin  
 Glycyl-L-proline-7-amido-4-methylcoumarin-hydrobromide  
 L-Histidine-7-amido-4-methylcoumarin  
 L-Isoleucine-7-amido-4-methylcoumarin  
 5 L-Isoleucine-7-amido-4-methylcoumarin TFA  
 L-Leucine-7-amido-4-methylcoumarin  
 L-Leucine-7-amido-4-methylcoumarin hydrochloride  
 L-Leucyl-L-valyl-L-tyrosine-7-amido-4-methylcoumarin  
 L-Lysine-7-amido-4-methylcoumarin acetate  
 10 L-Methionine-7-amido-4-methylcoumarin acetate  
 N-Methoxysuccinyl-L-alanyl-L-phenylalanyl-L-lysine-7-amido-4-methylcoumarin  
 TFA  
 N-Methoxysuccinyl-L-aspartyl-L-tyrosyl-L-methionine-7-amido-4-methylcoumarin  
 N-Methoxysuccinylglycyl-L-tryptophyl-L-methionine-7-amido-4-methylcoumarin  
 15 L-Ornithine-7-amido-4-methylcoumarin carbonate  
 L-Phenylalanine-7-amido-4-methylcoumarin TFA  
 L-Proline-7-amido-4-methylcoumarin hydrobromide  
 L-Prolyl-L-phenylalanyl-L-arginine-7-amido-4-methylcoumarin salt  
 L-Pyroglutamic acid-7-amido-4-methylcoumarin  
 20 L-Serine-7-amido-4-methylcoumarin hydrochloride  
 L-Seryl-L-tyrosine-7-amido-4-methylcoumarin Hydrate  
 N-Succinyl-L-alanyl-L-alanyl-L-alanine-7-amido-4-methylcoumarin  
 N-Succinyl-L-alanyl-L-alanyl-L-phenylalanine-7-amido-4-methylcoumarin  
 N-Succinyl-L-alanyl-L-alanyl-L-valine-7-amido-4-methylcoumarin  
 25 N-Succinyl-L-alanyl-L-phenylalanyl-L-lysine-7-amido-4-methylcoumarin  
 N-Succinyl-L-alanyl-L-phenylalanyl-L-lysine-7-amido-4-methylcoumarin TFA  
 N-Succinyl-L-alanyl-L-prolyl-L-alanine-7-amido-4-methylcoumarin  
 N-Succinylglycyl-L-proline-7-amido-4-methylcoumarin  
 N-p-Tosylglycyl-L-prolyl-L-arginine-7-amido-4-methylcoumarin hydrochloride  
 30 N-p-Tosylglycyl-L-prolyl-L-lysine-7-amido-4-methylcoumarin hydrochloride  
 L-Tyrosine-7-amido-4-methylcoumarin

Various Substrates

- L-alanine- $\beta$ -naphthylamide  
DL-Alanine- $\beta$ -naphthylamide hydrochloride  
L-Alanyl-L-alanine- $\beta$ -naphthylamide
- 5 p-Aminobenzyl-1-thio-2-acetamido-2-deoxy- $\beta$ -D-glucopyranoside  
p-Aminobenzyl-1-thio- $\beta$ -D-galactopyranoside  
D-Amygdalin from Apricot Kernels  
L-Arginine-4-methoxy- $\beta$ -naphthylamide hydrochloride  
L-Arginine- $\beta$ -naphthylamide hydrochloride
- 10 N-a-Benzoyl-L-arginine ethylester hydrochloride  
N-a-Benzoyl-L-arginine-4-methoxy- $\beta$ -naphthylamide hydrochloride  
N-a-Benzoyl-DL-arginine- $\beta$ -naphthylamide  
N-a-Benzoyl-D-arginine-p-nitroanalide hydrochloride  
N-a-Benzoyl-D-arginine-p-nitroanalide hydrochloride
- 15 N-a-Benzoyl-D-arginine-p-nitroanalide hydrochloride  
6-Benzoyl-2-naphthylphosphate Disodium salt  
6-Benzoyl-2-naphthylsulfate Potassium salt  
Bis(4-nitrophenyl) phosphate Sodium salt  
4-Bromomethyl-7-methoxycoumarin
- 20 6-Bromo-2-naphthyl acetate  
6-Bromo-2-naphthyl-N-acetyl- $\beta$ -D-glucosaminide  
6-Bromo-2-naphthyl- $\beta$ -D-galactoside  
6-Bromo-2-naphthyl- $\alpha$ -D-glucopyranoside  
6-Bromo-2-naphthyl- $\beta$ -D-glucopyranoside
- 25 6-Bromo-2-naphthyl- $\beta$ -D-glucuronide  
6-Bromo-2-naphthyl sulfate  
6-Bromo-2-naphthyl sulfate Potassium salt  
6-Bromo-2-naphthyl- $\beta$ -D-xylopyranoside  
2-Chloro-4-nitrophenyl-N-acetyl- $\beta$ -D-glucosaminide
- 30 2-Chloro-4-nitrophenyl- $\beta$ -D-cellobloside  
2-Chloro-4-nitrophenyl- $\beta$ -D-xylopyranoside  
8-Hydroxyquinoline- $\beta$ -D-glucuronide  
L-Leucine-p-nitroanilide  
L-Leucyl-4-methoxy- $\beta$ -naphthylamide
- 35 L-Leucyl- $\beta$ -naphthylamide  
DL-Methionine- $\beta$ -naphthylamide hydrochloride  
2-(3'-Methoxyphenyl)-N-acetyl-D-neuraminic acid  
Naphthol AS  
Naphthol AS-acetate
- 40 Naphthol AS-B1-N-acetyl- $\beta$ -D-glucosaminide  
Naphthol AS- $\beta$ -chloropropionate  
Naphthol AS-B1- $\beta$ -L-fucopyranoside  
Naphthol AS-B1- $\beta$ -D-galactopyranoside  
Naphthol AS-B1- $\beta$ -D-galactosaminide
- 45 Naphthol AS-B1-glucopyranoside  
Naphthol AS-B1- $\beta$ -D-glucuronic acid  
Naphthol AS-nonanoate

Table I

- Naphthol AS- $\gamma$ -phenylbutyrate  
Naphthol AS-phenylpropionate  
Naphthol AS-phosphate  
Naphthol AS-B1-phosphate  
5 Naphthol AS-phosphate Sodium salt  
Naphthol AS-B1-phosphate Sodium salt  
Naphthol AS-sulphate Potassium salt  
Naphthol AS-B1-sulfate Potassium salt  
1-Naphthylbutyrate  
10 2-Naphthylbutyrate  
1-Naphthylcaprylate  
2-Naphthylcaprylate  
1-Naphthyl- $\alpha$ -D-galactopyranoside  
1-Naphthyl- $\beta$ -D-galactopyranoside  
15 1-Naphthyl- $\beta$ -D-galactopyranoside  
1-Naphthyl- $\beta$ -D-glucuronide  
1-Naphthylphosphate Disodium salt  
2-Naphthylphosphate Disodium salt  
2-Naphthylphosphate Sodium salt  
20 2-Naphthylphosphate Sodium salt  
1-Naphthylphosphate Sodium salt  
2-Naphthylsulfate Potassium salt  
2-Nitrophenyl-acetate  
4-Nitrophenyl-acetate  
25 2-Nitrophenyl-N-acetyl- $\alpha$ -D-galactosaminide  
4-Nitrophenyl-N-acetyl- $\alpha$ -D-galactosaminide  
4-Nitrophenyl-N-acetyl- $\beta$ -D-galactosaminide  
4-Nitrophenyl-N-acetyl- $\alpha$ -D-glucosaminide  
4-Nitrophenyl-N-acetyl- $\beta$ -D-glucosaminide  
30 4-Nitrophenyl-N-acetyl-1-thio- $\beta$ -D-glucosaminide  
4-Nitrophenyl- $\alpha$ -L-arabinopyranoside  
2-Nitrophenyl-butyrate  
4-Nitrophenyl-butyrate  
4-Nitrophenyl-caprate  
35 4-Nitrophenyl-caproate  
3-Nitrophenyl-caprylate  
4-Nitrophenyl-caprylate  
4-Nitrophenyl- $\beta$ -D-cellobloside  
3-Nitrophenyl- $\beta$ -D-fucopyranoside  
40 4-Nitrophenyl- $\alpha$ -D-fucopyranoside  
4-Nitrophenyl- $\beta$ -D-fucopyranoside  
4-Nitrophenyl- $\alpha$ -L-fucopyranoside  
4-Nitrophenyl- $\beta$ -L-fucopyranoside  
2-Nitrophenyl- $\alpha$ -D-galactopyranoside  
45 2-Nitrophenyl- $\beta$ -D-galactopyranoside  
3-Nitrophenyl- $\alpha$ -D-galactopyranoside  
3-Nitrophenyl- $\beta$ -D-galactopyranoside



- 4-Nitrophenyl- $\alpha$ -D-galactopyranoside
- 4-Nitrophenyl- $\beta$ -D-galactopyranoside
- 2-Nitrophenyl- $\beta$ -D-galactopyranoside-6-phosphate Cyclohexylammonium salt
- 4-Nitrophenyl- $\beta$ -D-galacturonide
- 5 4-Nitrophenyl- $\alpha$ -D-glucopyranoside
- 4-Nitrophenyl- $\beta$ -D-glucopyranoside
- 4-Nitrophenyl- $\beta$ -D-glucuronide
- 2-Nitrophenyl- $\beta$ -D-glucuronide
- 4-Nitrophenyl-glycerol
- 10 4-Nitrophenyl-4'-guanidinobenzoate
- 4-Nitrophenyl- $\alpha$ -D-maltoheptaoside
- 4-Nitrophenyl- $\alpha$ -D-maltohexaoside
- 4-Nitrophenyl- $\alpha$ -D-maltopentaoside
- 4-Nitrophenyl- $\alpha$ -D-maltoside
- 15 4-Nitrophenyl- $\alpha$ -D-maltatetraoside
- 4-Nitrophenyl- $\alpha$ -D-maltatrioside
- 4-Nitrophenyl- $\alpha$ -D-mannopyranoside
- 4-Nitrophenyl- $\beta$ -D-mannopyranoside
- 2-Nitrophenyl-myristate
- 20 4-Nitrophenyl-myristate
- 2-Nitrophenyl-palmitate
- 4-Nitrophenyl-palmitate
- p-Nitrophenylphosphate Disodium salt Hexahydrate high purity
- 4-Nitrophenyl-proionate
- 25 4-Nitrophenyl-stearate
- 4-Nitrophenyl-sulfate Potassium salt
- 2-Nitrophenyl- $\beta$ -D-thiogalactopyranoside
- 4-Nitrophenyl- $\beta$ -D-thiogalactopyranoside
- 4-Nitrophenyl- $\beta$ -D-thioglucopyranoside
- 30 4-Nitrophenyl- $\beta$ -D-xylopyranoside
- Phenolphthalein diphosphate
- Phenolphthalein diphosphate Tetrasodium salt
- Phenolphthalein-mono- $\beta$ -D-galactopyranoside
- Phenolphthalein- $\beta$ -D-glucuronic acid Sodium salt
- 35 Phenyl-N-acetyl- $\alpha$ -D-glucosaminide
- Phenylethyl- $\beta$ -D-galactoside
- Phenyl- $\beta$ -D-galactoside
- Phenyl- $\alpha$ -D-glucoside
- Phenyl- $\alpha$ -D-glucoside tetraacetate
- 40 Phenyl- $\beta$ -D-glucoside tetraacetate
- Resorufin- $\beta$ -D-galactopyranoside
- Resorufin- $\beta$ -D-glucuronide
- L-Serine- $\beta$ -naphthylamide
- 1-Thio- $\beta$ -D-galactopyranoside Sodium salt
- 45 1-Thio- $\beta$ -D-glucopyranoside Sodium salt
- L-Tyrosine- $\beta$ -naphthylamide

X-SUBSTRATES (5-Bromo-4-chloro-3-Indolyl-Substrates)

- 5-Bromo-4-chloro-3-indolyl-acetate  
 5-Bromo-4-chloro-3-indolyl-N-acetyl- $\beta$ -D-galactosaminide  
 5-Bromo-4-chloro-3-indolyl-N-acetyl- $\beta$ -D-glucosaminide  
 5 5-Bromo-4-chloro-3-indolyl-butyrate  
 5-Bromo-4-chloro-3-indolyl-caprylate  
 5-Bromo-4-chloro-3-indolyl-Carbohydrates and other Derivates  
 5-Bromo-4-chloro-3-indolyl-1,3-diacetate  
 5-Bromo-4-chloro-3-indolyl- $\beta$ -D-fucopyranoside  
 10 5-Bromo-4-chloro-3-indolyl- $\alpha$ -D-galactopyranoside  
 5-Bromo-4-chloro-3-indolyl- $\beta$ -D-glucopyranoside  
 5-Bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronic acid Cyclohexylammonium salt  
 5-Bromo-4-chloro-3-indolyl- $\beta$ -D-glucuronic acid Sodium salt  
 5-Bromo-4-chloro-3-indolyl- $\alpha$ -D-mannopyranoside  
 15 5-Bromo-4-chloro-3-indolyl-phosphate Disodium salt  
 5-Bromo-4-chloro-3-indolyl-phosphate Potassium salt  
 5-Bromo-4-chloro-3-indolyl-phosphate p-Toluidine salt  
 5-Bromo-4-chloro-3-indolyl-sulfate Potassium salt  
 5-Bromo-4-chloro-3-indolyl- $\beta$ -D-xylopyranoside

20 Y-SUBSTRATES (Indoxyl-Substrates)

- 8-Bromoindoxyl-3-acetate  
 5-Bromoindoxyl-1,3-diacetate  
 Indoxyl-1,3-diacetate  
 Indoxyl- $\beta$ -D-galactoside  
 25 Indoxyl- $\beta$ -D-glucoside  
 Indoxyl- $\beta$ -D-glucuronic acid Cyclohexylammonium salt  
 3-Indoxyl-phosphate Di(2-amino-2-methyl-1,3-propanediol) salt  
 3-Indoxyl-phosphate Disodium salt  
 3-Indoxyl-phosphate p-Toluidine salt  
 30 3-Indoxylsulfate Potassium salt

Table II: Media Formulation (per liter)

		(grams)
	Defined media	15.36
	HEPES (acid)	4.29
5	HEPES (Na <sup>+</sup> salt)	8.38
	Bacto Proteose peptone No. 3 (Difco)	5.00
	Potassium nitrate	5.00
	4-methylumbelliferyl phosphate (Sigma)	0.025
	4-methylumbelliferyl- $\beta$ -D-glucoside (Sigma)	0.025
10	L-alanine-7-amido-4-methyl coumarin (Sigma)	0.025

Table III: Defined Media Composition

INGREDIENT		CONCENTRATION (mg/L)
	Ammonium acetate	500
	Magnesium chloride	95.35
5	Ferric chloride (6 hydrate)	2.7
	Manganese sulfate (1 hydrate)	0.273
	Potassium chloride	100
	Zinc sulfate (7 hydrate)	0.8
	Calcium chloride (2 hydrate)	7.38
10	Sodium chloride	1000
	L-arginine HCl	1270
	L-asparagine (1 hydrate)	1136
	L-aspartic acid	20
	L-cysteine HCl (1 hydrate)	1450
15	L-cystine methylester 2 HCl	340.8
	L-glutamic acid	20
	L-glutamine	2520
	Glycine	500
	L-histidine HCl (1 hydrate)	419
20	L-Isoleucine	520
	L-leucine	520
	L-lycine HCl	724.65
	L-methionine	150
	L-phenylalanine	320
25	L-proline	1000
	L-serine	30
	L-threonine	480
	L-tryptophan	100
	L-tyrosine Na salt (2 hydrate)	519
30	L-valine	460
	Adenine	25
	Biotin	0.5
	Choline chloride	25
	Folic acid	5
35	I-Inositol	25
	D(+)calcium pantothenate	25
	Nicotinamide	5
	Para aminobenzoic acid	1
	Pyridoxal HCl	5
40	Riboflavin	5
	Thiamine HCl	5
	Uracil	25
	Sodium pyruvate	1000

What is Claimed is:

1. Method for detecting the existence or measuring the concentration of bacteria in a test sample, comprising the steps of:

5 providing a bacterial growth medium comprising three or more different enzyme substrates, wherein each said substrate is hydrolysed by a different bacterial enzyme, and thereafter, causes or produces a detectable signal;

10 inoculating said medium with said test sample and incubating said medium under a condition suitable for bacterial growth for a certain time period; and

detecting or measuring the detectable signal as an indication of the existence or the concentration of bacteria in said test sample.

15 2. The method of claim 1, wherein said different substrates each having both a nutrient moiety and a detectable moiety linked together by a covalent bond, and each said substrate is hydrolysed by a different bacterial enzyme to produce a separate detectable moiety, and said  
20 separate detectable moiety causes or produces a detectable signal.

3. The method of claim 1, wherein said bacteria are selected from the group consisting of *Aeromonas hydrophilia*, *Aeromonas caviae*, *Aeromonas sobria*,  
25 *Streptococcus uberis*, *Enterococcus faecium*, *Enterococcus faecalis*, *Bacillus sphaericus*, *Pseudomonas fluorescens*, *Pseudomonas putida*, *Serratia liquefaciens*, *Lactococcus lactis*, *Xanthomonas maltophilia*, *Staphylococcus simulans*, *Staphylococcus hominis*, *Streptococcus constellatus*,  
30 *Streptococcus anginosus*, *Escherichia coli*, *Staphylococcus aureus*, *Mycobacterium fortuitum*, and *Klebsiella pneumonia*.

4. The method of claim 1, wherein said bacterial enzyme is selected from the group consisting of alkaline phosphatase, acid phosphatase, esterase, lipase, N-acetyl-

$\beta$ -D-galactosaminidase, N-acetyl- $\beta$ -D-glucosaminidase, Neuraminidase, L-arabinopyranosidase,  $\beta$ -D-fucosidase,  $\alpha$ -L-fucosidase,  $\beta$ -L-fucosidase,  $\alpha$ -D-galactosidase,  $\beta$ -D-galactosidase,  $\alpha$ -D-glucosidase,  $\beta$ -D-glucosidase,  $\beta$ -D-glucuronidase,  $\alpha$ -D-mannosidase, pyrophosphatase, sulfatase,  $\beta$ -D-xylosidase, peptidase, (L or D amino acid) - aminopeptidase, L-alanine aminopeptidase, trypsin, chymotrypsin, and phosphohydrolase.

5. The method of claim 1, wherein one of said  
10 substrates is hydrolysed by a phosphatase enzyme, another of said substrates is hydrolysed by a glycosidase enzyme, and a third said substrate is hydrolysed by a peptidase enzyme.

6. The method of claim 1, wherein said detectable  
15 moiety is a fluorescent moiety and said detectable signal is a fluorescent signal.

7. The method of claim 1, wherein said substrates  
comprise 4-methylumbelliferyl phosphate, 4-methylumbelliferyl- $\beta$ -D-glucoside and L-alanine-7-amido-4-methyl  
20 coumarin.

8. The method of claim 1, wherein said test sample  
is taken from a food product.

9. The method of claim 8, wherein said food product  
is ground beef.

25 10. The method of claim 8, wherein said food product  
is chicken.

11. The method of claim 8, wherein said food product  
is water.

12. The method of claim 1, wherein said medium is liquid.

13. The method of claim 1, wherein said time period is no more than 24 hours.

5 14. Method for detecting the existence or measuring the concentration of bacteria in a test sample, comprising the steps of:

providing a bacterial growth medium comprising two or more different substrates, wherein each said substrate is hydrolysed by a different bacterial enzyme and thereafter causes or produces an identical type of detectable signal;

10 inoculating said medium with said test sample and incubating said medium under a condition suitable for bacterial growth for a certain time period; and

15 detecting or measuring the detectable signal as an indication of the existence or the concentration of bacteria in said test sample.

15. The method of claim 14, wherein said different substrates each having both a nutrient moiety and a detectable moiety linked together by a covalent bond, and each said substrate is hydrolysed by a different bacterial enzyme to produce a separate detectable moiety, and said separate detectable moiety causes or produces an identical type of detectable signal.

25 16. The method of claim 14, wherein said bacteria are selected from the group consisting of *Aeromonas hydrophilia*, *Aeromonas caviae*, *Aeromonas sobria*, *Streptococcus uberis*, *Enterococcus faecium*, *Enterococcus faecalis*, *Bacillus sphaericus*, *Pseudomonas fluorescens*,  
30 *Pseudomonas putida*, *Serratia liquefaciens*, *Lactococcus lactis*, *Xanthomonas maltophilia*, *Staphylococcus simulans*, *Staphylococcus hominis*, *Streptococcus constellatus*,

*Streptococcus anginosus*, *Escherichia coli*, *Staphylococcus aureus*, *Mycobacterium fortuitum*, and *Klebsiella pneumonia*.

17. The method of claim 14, wherein said bacterial enzyme is selected from the group consisting of alkaline  
5 phosphatase, acid phosphatase, esterase, lipase, N-acetyl- $\beta$ -D-galactosaminidase, N-acetyl- $\beta$ -D-glucosaminidase, Neuraminidase, L-arabinopyranosidase,  $\beta$ -D-fucosidase,  $\alpha$ -L-fucosidase,  $\beta$ -L-fucosidase,  $\alpha$ -D-galactosidase,  $\beta$ -D-galactosidase,  $\alpha$ -D-glucosidase,  $\beta$ -D-glucosidase,  $\beta$ -D-  
10 glucuronidase,  $\alpha$ -D-mannosidase, pyrophosphatase, sulfatase,  $\beta$ -D-xylosidase, peptidase, (L or D amino acid)-aminopeptidase, L-alanine aminopeptidase, trypsin, chymotrypsin, and phosphohydrolase.

18. The method of claim 14, wherein said enzyme is  
15 selected from the group consisting of a phosphatase enzyme, a glycosidase enzyme and a peptidase enzyme.

19. The method of claim 14, wherein said detectable moiety is a fluorescent moiety and said detectable signal is a fluorescent signal.

20. The method of claim 18 or 19, wherein said  
20 substrates are selected from the group consisting of 4-methylumbelliferyl phosphate, 4-methylumbelliferyl- $\beta$ -D-glucoside and L-alanine-7-amido-4-methyl coumarin.

21. The method of claim 14, wherein said test sample  
25 is taken from a food product.

22. The method of claim 21, wherein said food product is ground beef.

23. The method of claim 21, wherein said food product is chicken.



24. The method of claim 21, wherein said food product is water.

25. The method of claim 14, wherein said medium is liquid.

5 26. The method of claim 14, wherein said time period is no more than 24 hours.

27. A bacterial growth medium comprising three or more different substrates, wherein each said substrate is hydrolysed by a different bacterial enzyme, and thereafter, causes or produces a detectable signal.  
10

28. The medium of claim 27, wherein said different substrates each having both a nutrient moiety and a detectable moiety linked together by a covalent bond, and each said substrate is hydrolysed by a different bacterial enzyme to produce a separate detectable moiety, and said separate detectable moiety causes or produces a detectable signal.  
15

29. The medium of claim 27, wherein said bacterial enzyme is selected from the group consisting of alkaline phosphatase, acid phosphatase, esterase, lipase, N-acetyl- $\beta$ -D-galactosaminidase, N-acetyl- $\beta$ -D-glucosaminidase, Neuraminidase, L-arabinopyranosidase,  $\beta$ -D-fucosidase,  $\alpha$ -L-fucosidase,  $\beta$ -L-fucosidase,  $\alpha$ -D-galactosidase,  $\beta$ -D-galactosidase,  $\alpha$ -D-glucosidase,  $\beta$ -D-glucosidase,  $\beta$ -D-glucuronidase,  $\alpha$ -D-mannosidase, pyrophosphatase, sulfatase,  $\beta$ -D-xylosidase, peptidase, (L or D amino acid) - aminopeptidase, L-alanine aminopeptidase, trypsin, chymotrypsin, and phosphohydrolase.  
20  
25

30. The medium of claim 27, wherein one of said substrates is hydrolysed by a phosphatase enzyme, another of said substrates is hydrolysed by a glycosidase enzyme,  
30

and a third said substrate is hydrolysed by a peptidase enzyme.

31. The medium of claim 27, wherein said detectable moiety is a fluorescent moiety and said detectable signal  
5 is a fluorescent signal.

32. The medium of claim 27, wherein said substrates comprise 4-methylumbelliferyl phosphate, 4-methylumbelliferyl- $\beta$ -D-glucoside and L-alanine-7-amido-4-methyl coumarin.

10 33. The medium of claim 27, further comprising a test sample from a food product.

34. The medium of claim 33, wherein said food product is ground beef.

15 35. The medium of claim 33, wherein said food product is chicken.

36. The medium of claim 33, wherein said food product is water.

37. The medium of claim 27, wherein said medium is liquid.

20 38. A bacterial growth medium comprising two or more different substrates, wherein each said substrate is hydrolysed by a different bacterial enzyme and thereafter causes or produces an identical type of detectable signal.

25 39. The medium of claim 38, wherein said different substrates each having both a nutrient moiety and a detectable moiety linked together by a covalent bond, and each said substrate is hydrolysed by a different bacterial enzyme to produce a separate detectable moiety, and said

separate detectable moiety causes or produces an identical type of detectable signal.

40. The medium of claim 38, wherein said bacterial enzyme is selected from the group consisting of alkaline phosphatase, acid phosphatase, esterase, lipase, N-acetyl- $\beta$ -D-galactosaminidase, N-acetyl- $\beta$ -D-glucosaminidase, Neuraminidase, L-arabinopyranosidase,  $\beta$ -D-fucosidase,  $\alpha$ -L-fucosidase,  $\beta$ -L-fucosidase,  $\alpha$ -D-galactosidase,  $\beta$ -D-galactosidase,  $\alpha$ -D-glucosidase,  $\beta$ -D-glucosidase,  $\beta$ -D-glucuronidase,  $\alpha$ -D-mannosidase, pyrophosphatase, sulfatase,  $\beta$ -D-xylosidase, peptidase, (L or D amino acid) - aminopeptidase, L-alanine aminopeptidase, trypsin, chymotrypsin, and phosphohydrolase.

41. The medium of claim 38, wherein said enzyme is selected from the group consisting of a phosphatase enzyme, a glycosidase enzyme and a peptidase enzyme.

42. The medium of claim 38, wherein said detectable moiety is a fluorescent moiety and said detectable signal is a fluorescent signal.

43. The medium of claim 41 or 42, wherein said substrates are selected from the group consisting of 4-methylumbelliferyl phosphate, 4-methylumbelliferyl- $\beta$ -D-glucoside and L-alanine-7-amido-4-methyl coumarin.

44. The medium of claim 38, further comprising a test sample from a food product.

45. The medium of claim 44, wherein said food product is ground beef.

46. The medium of claim 44, wherein said food product is chicken.

47. The medium of claim 44, wherein said food product is water.

48. The medium of claim 38, wherein said medium is liquid.

5 49. Method for detecting the existence or measuring the concentration of eukaryotic microbes in a test sample, comprising the steps of:

providing a growth medium comprising three or more different substrates, wherein each said substrate is  
10 hydrolysed by a different eukaryotic microbial enzyme and thereafter causes or produces a detectable signal;

inoculating said medium with said test sample and incubating said medium under a condition suitable for microbial growth for a certain time period; and

15 detecting or measuring the detectable signal as an indication of the existence or the concentration of eukaryotic microbes in said test sample.

50. The method of claim 49, wherein said different substrates each having both a nutrient moiety and a  
20 detectable moiety linked together by a covalent bond, and each said substrate is hydrolysed by a different eukaryotic microbial enzyme to produce a separate detectable moiety, and said separate detectable moiety causes or produces a detectable signal.

25 51. The method of claim 49, wherein said eukaryotic microbes comprise a yeast.

52. The method of claim 5 or 18, wherein said peptidase enzyme is an aminopeptidase enzyme.

53. The medium of claim 30 or 41, wherein said peptidase enzyme is an aminopeptidase enzyme.  
30

## INTERNATIONAL SEARCH REPORT

International Application No.

PC, /US 96/08124

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12Q1/04 G01N33/02 G01N33/18

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	US,A,5 464 755 (BOCHNER BARRY) 7 November 1995  see claim 1 ---	1-4, 12-18, 26-30, 37-41,48
E	FR,A,2 728 587 (PASTEUR SANOFI DIAGNOSTICS) 28 June 1996 see the whole document ---	14-18, 38-41
P,X	US,A,5 443 987 (DECICCO BENEDICT T ET AL) 22 August 1995  see claims 6,7 --- -/--	14-18, 25,26, 38-41,48



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*&\* document member of the same patent family

Date of the actual completion of the international search

18 September 1996

Date of mailing of the international search report

08.10.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Wells, A

## INTERNATIONAL SEARCH REPORT

International Application No

PC, US 96/08124

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP,A,0 122 028 (E Y LAB INC) 17 October 1984	1-5, 12-18, 25-30, 38-41, 47,48, 52,53
Y	see page 4 - page 6; example 3	6-11,22, 23,32, 34,45, 46,49-51
X	WO,A,95 04157 (RAMBACH ALAIN) 9 February 1995	14-18, 21, 24-26, 38-41, 44,47,48
Y	see page 1, line 4-5; tables II,III	8-11,22, 23,34, 45,46, 49-51
X	EP,A,0 254 771 (EDBERG STEPHEN C) 3 February 1988	14-21, 24-29, 31,33, 36-44, 47,48
Y	see page 8, line 17-25 - page 9	6-11,22, 23,32, 34,45,46
X	US,A,5 364 767 (FLOWERS DANIEL G ET AL) 15 November 1994 see column 13	14-18, 25,26, 38-41,48

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC, /US 96/08124

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-5464755	07-11-95	AU-A- 2706295 WO-A- 9529984 US-A- 5541082	29-11-95 09-11-95 30-07-96
FR-A-2728587	28-06-96	NONE	
US-A-5443987	22-08-95	NONE	
EP-A-0122028	17-10-84	JP-A- 59220199 US-A- 5089394	11-12-84 18-02-92
WO-A-9504157	09-02-95	FR-A- 2708286 CA-A- 2168114 EP-A- 0711360	03-02-95 09-02-95 15-05-96
EP-A-0254771	03-02-88	US-A- 4925789 AU-B- 597675 AU-A- 7479587 CA-A- 1288322 DE-A- 3687232 ES-T- 2044835 IE-B- 60884 JP-B- 7004272 JP-A- 63014699 US-A- 5429933	15-05-90 07-06-90 07-01-88 03-09-91 14-01-93 16-01-94 24-08-94 25-01-95 21-01-88 04-07-95
US-A-5364767	15-11-94	NONE	